

## Questions and Answers

### **Recommended composition of influenza virus vaccines for use in the 2018-2019 northern hemisphere influenza season and development of candidate vaccine viruses for pandemic preparedness**

22 February 2018

1. What is the WHO Global Influenza Surveillance and Response System (GISRS)?
2. What is the purpose of WHO recommendations on the composition of influenza virus vaccines?
3. How are influenza vaccine recommendations made?

oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo

4. What viruses are recommended by WHO to be included in influenza vaccines for use in the 2018-2019 northern hemisphere influenza season?
5. Are the vaccine viruses in this recommendation different from those in previous recommendations?
6. Why was the H3 component of the vaccine for the 2018-19 northern hemisphere influenza season changed to A/Singapore/INFIMH-16-0019/2016-like virus?
7. What is the difference between trivalent and quadrivalent vaccines?
8. Why was the influenza B/Victoria lineage virus component for vaccines changed from the B/Brisbane/60/2008-like virus to a B/Colorado/06/2017-like virus?
9. What are the implications of changing the influenza B component of the northern hemisphere vaccines for the upcoming southern hemisphere influenza season where the B/Phuket/3073/2013-like virus (B/Yamagata lineage) is in the trivalent vaccines and B/Brisbane/60/2008-like virus (B/Victoria lineage) is in the quadrivalent vaccines?

oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo oo

10. What vaccine formulation (i.e. recommendation for northern or southern hemisphere influenza season) should countries in tropical and subtropical regions consider for use in vaccination campaigns?
11. What are candidate vaccine viruses (CVVs)?
12. What CVVs are available for use in influenza vaccines?
13. What happens after the WHO recommendations are made?
14. Why does GISRS continue to update the list of available CVVs for pandemic preparedness?

## **1. What is the WHO Global Influenza Surveillance and Response System (GISRS)?**

GISRS is a global public health laboratory network coordinated by WHO, currently consisting of 144 National Influenza Centres (NICs) in 114 WHO Member States, 6 WHO Collaborating Centres for Influenza (CCs), 4 WHO Essential Regulatory Laboratories (ERLs) and 13 WHO H5 Reference Laboratories.

This network monitors evolution of influenza viruses of public health concern, such as seasonal, zoonotic and pandemic viruses, conducts risk assessment, recommends and implements risk response measures. In the last year, NICs collected and tested more than three million clinical specimens from patients and shared representative influenza viruses with the WHO CCs for detailed analyses and for making recommendations for vaccine composition. The network also provides guidance to countries and support for activities such as training, risk assessment, outbreak response, development of diagnostic tests, testing for antiviral drug resistance and scientific interpretation of important findings.

## **2. What is the purpose of WHO recommendations on the composition of influenza virus vaccines?**

WHO recommendations provide a guide to national public health and regulatory authorities and vaccine manufacturers for the development and production of influenza vaccines for the next influenza season. In contrast to many other vaccines, the viruses in influenza vaccines have to be evaluated and updated regularly because circulating influenza viruses continuously evolve. Recommendations are made in February/March for the following influenza season in the northern hemisphere and in September for the following influenza season in the southern hemisphere because approximately 6-8 months are needed to produce and approve vaccines.

## **3. How are influenza vaccine recommendations made?**

Many different sources of data and information are used to determine the recommended vaccine viruses, including:

- ***Surveillance data from the GISRS network, which includes NICs, WHO CCs, WHO ERLs and WHO H5 Reference Laboratories:***  
Virus surveillance data, complemented with epidemiologic and clinical findings inform the vaccine virus selection process.
- ***Antigenic characterization of viruses:***  
GISRS laboratories, in particular the WHO CCs, also conduct testing to evaluate the antibody or immune response triggered by the proteins on the surface of influenza viruses. Antigenic cartography is used as a way to visualize relatedness of viruses.
- ***Human serology studies with inactivated influenza virus vaccines:***  
WHO CCs and WHO ERLs use tests to determine how well antibodies from vaccinated people react with recently circulating influenza viruses.
- ***Genetic characterization of viruses:***  
GISRS laboratories conduct testing to compare virus gene sequences of circulating

influenza viruses to the sequences of vaccine viruses to identify genetic changes that might influence protection conferred by a given vaccine.

- ***Virus fitness forecasting:***  
Information from modelling studies, based on genetic and antigenic information, is also considered.
- ***Antiviral resistance:***  
GISRS laboratories test influenza viruses to determine if they have any resistance to the antiviral drugs used to treat influenza infection. This information is taken into consideration when specific viruses are selected as CVVs.
- ***Vaccine effectiveness:***  
The Global Influenza Vaccine Effectiveness (GIVE) Collaboration, made up of 19 different studies conducted in countries in both the northern and southern hemispheres, provides information on vaccine performance in previous and current influenza seasons.
- ***Availability of potential CVVs:***  
The vast majority of vaccines produced globally use egg-based manufacturing processes. This requires CVVs which grow well in eggs. These viruses must be available in order to produce vaccine and make the vaccine available in time for the next influenza season.

These data, and other findings made available by GISRS laboratories, are evaluated during WHO Consultations in February/March and September of each year. The consultation includes Advisers from WHO CCs and WHO ERLs, as well as Observers and other experts from WHO CCs, WHO ERLs, WHO H5 Reference Laboratories, NICs, the University of Cambridge, the OIE/FAO Network of expertise on animal influenza (OFFLU), and other national and regional institutions. Further information about GISRS is available at [http://www.who.int/influenza/gisrs\\_laboratory/en/](http://www.who.int/influenza/gisrs_laboratory/en/).

#### **4. What viruses are recommended by WHO to be included in influenza vaccines for use in the 2018-2019 northern hemisphere influenza season?**

WHO recommends that quadrivalent influenza vaccines for use in the 2018-19 northern hemisphere influenza season contain the following viruses:

- an A/Michigan/45/2015 (H1N1)pdm09-like virus;
- an A/Singapore/INFIMH-16-0019/2016 (H3N2)-like virus;
- a B/Colorado/06/2017-like virus; and
- a B/Phuket/3073/2013-like virus

It is recommended that trivalent vaccines contain the B/Colorado/06/2017-like virus, of the B/Victoria/2/87-lineage.

**5. Are the vaccine viruses in this recommendation different from those in previous recommendations?**

The influenza B/Victoria lineage virus component and the A(H3N2) virus component have been updated compared to the viruses recommended for the 2017-2018 northern hemisphere influenza seasons. The update is as follows:

- replacement of the A/Hong Kong/4801/2014 (H3N2)-like virus with an A/Singapore/INFIMH-16-0019/2016 (H3N2)-like virus.
- replacement of the B/Brisbane/60/2008-like virus with a B/Colorado/06/2017-like virus.

All previous WHO recommendations can be found on the WHO Global Influenza Programme website at: <http://www.who.int/influenza/vaccines/virus/recommendations/en/>

**6. Why was the H3 component of the vaccine for the 2018-19 northern hemisphere influenza season changed to A/Singapore/INFIMH-16-0019/2016-like virus?**

Circulating A(H3N2) viruses were poorly recognized by antibodies raised against egg-propagated A/Hong Kong/4801/2014 (H3N2)-like virus, the vaccine virus for the northern hemisphere 2017-2018. Recent A(H3N2) viruses were better inhibited by ferret antisera raised against the egg-propagated reference virus A/Singapore/INFIMH-16-0019/2016 compared to ferret antisera raised against other recent egg-propagated A(H3N2) viruses.

**7. What is the difference between quadrivalent and trivalent vaccines?**

Quadrivalent vaccines include two type A viruses (an A(H3N2) and an A(H1N1)pdm09) and two type B viruses (one from each lineage). Trivalent vaccines include three components, two type A viruses (an A(H3N2) and an A(H1N1)pdm09) and one type B virus.

**8. Why was the influenza B/Victoria lineage virus component for vaccines changed from the B/Brisbane/60/2008-like virus to a B/Colorado/06/2017-like virus?**

While most recent B/Victoria/2/87 lineage viruses were well inhibited by antisera raised against B/Brisbane/60/2008-like viruses, a substantial and increasing proportion of viruses were antigenically distinguishable from B/Brisbane/60/2008-like viruses, but closely related to B/Colorado/06/2017. The significant majority of these viruses had a two amino acid deletion in their haemagglutinin.

**9. Is the vaccine composition for the 2018 southern hemisphere influenza season still appropriate?**

Of the influenza B viruses, a B/Yamagata/16/88 lineage virus was recommended for trivalent vaccines for the 2018 southern hemisphere influenza season. So far B/Yamagata lineage viruses predominated, with small number of B/Victoria/2/87 lineage detected. Of the

B/Victoria lineage viruses, an increasing proportion of viruses with HA deletions of two amino acids have been detected in many countries, leading to a change in the recommendation for the B/Victoria lineage component and the virus recommended for use in the 2018-2019 northern hemisphere trivalent vaccine. Current data indicate that the deletion variant is not circulating widely in the southern hemisphere.

**10. What vaccine formulation (i.e. recommendation for northern or southern hemisphere influenza season) should countries in tropical and subtropical regions consider for use in vaccines?**

Influenza viruses circulate at varying times through the year in tropical and sub-tropical countries. In selecting which vaccine formulation to use, these countries should consider their surveillance information, in particular epidemiological and virological data to decide when to start vaccination and whether to use the formulation recommended for the northern or southern hemisphere influenza season. WHO has formulated guidance for countries in tropical and sub-tropical regions to assist them in choosing which vaccine composition (February/March or September) is most appropriate (<http://www.who.int/influenza/vaccines/tropics/en/>).

**11. What are candidate vaccine viruses (CVVs)?**

A CVV is a virus prepared for potential use in vaccine manufacturing that is antigenically similar to the virus that has been recommended for use in egg-based or cell culture-based vaccines.

**12. What CVVs are available for use in influenza vaccines?**

The WHO recommended CVVs for vaccine development and production for the 2018-19 northern hemisphere influenza season are listed at:

[http://www.who.int/influenza/vaccines/virus/candidates\\_reagents/2018\\_19\\_north/en/](http://www.who.int/influenza/vaccines/virus/candidates_reagents/2018_19_north/en/)

The availability of CVVs by type/subtype, including zoonotic viruses, and corresponding potency test reagents is posted and updated on the WHO web site:

[http://www.who.int/influenza/vaccines/virus/candidates\\_reagents/home/en/](http://www.who.int/influenza/vaccines/virus/candidates_reagents/home/en/)

**13. What happens after the WHO recommendations are made?**

Approval of the composition and formulation of vaccines that will be used in each country is the responsibility of national or regional regulatory authorities. It is the responsibility of the vaccine manufacturer to obtain the appropriate CVVs and to obtain approval from the local regulatory agency. WHO publishes and updates a list of CVVs for selection by the manufacturers and regulatory agencies.

([http://www.who.int/influenza/vaccines/virus/candidates\\_reagents/home/en/](http://www.who.int/influenza/vaccines/virus/candidates_reagents/home/en/))

#### **14. Why does GISRS continue to update the list of available CVVs for pandemic preparedness?**

Influenza viruses circulate widely in some animals and may transmit sporadically to humans, resulting in zoonotic infections. As part of an influenza pandemic preparedness program, the WHO GISRS in collaboration with animal health partners analyses a range of zoonotic and potentially pandemic influenza viruses as they emerge and evolve, and develops relevant CVVs as a first step in the production of influenza vaccines. The selection and development of a zoonotic CVV is done to maintain a bank of viruses suitable for the immediate development of vaccines, for example during a pandemic, and also to assist those who may want to make pilot lots of vaccines, conduct clinical trials, or perform other pandemic preparedness tasks. The decision to use these materials for vaccine development should be based on an assessment of the public health risk and needs in consultation with national regulatory and public health authorities.

Further information about zoonotic influenza CVVs can be found at:

[http://www.who.int/influenza/vaccines/virus/characteristics\\_virus\\_vaccines/en/](http://www.who.int/influenza/vaccines/virus/characteristics_virus_vaccines/en/)

**For more information, please contact the WHO Global Influenza Programme at [GISRS-whohq@who.int](mailto:GISRS-whohq@who.int)**