INTERAGENCY INTEGRATED TRIAGE TOOL: Age < 12



(1)

CHECK FOR RED CRITERIA

Unresponsive

AIRWAY & BREATHING

- Stridor
- Respiratory distress* or central cyanosis

CIRCULATION

- Capillary refill >3 sec
- Weak and fast pulse
- Heavy bleeding
- Cold extremities
- Any two of:
- Letharqy
- Sunken eyes
- Very slow skin pinch
- Drinks poorly

DISABILITY

- Active convulsions
- Altered mental status (confused, restless, continuously irritable or lethargic) with stiff neck, hypothermia or fever
- Hypoglycaemia (if known)

OTHER MANAGEMENT

- Any infant <8 days old
- Age <2 months and temp <36 or >39°C
- High-risk trauma*
- Threatened limb*
- Acute testicular/scrotal pain or priapism
- Snake bite
- Poisoning/ingestion or dangerous chemical exposure*
- Pregnant with adult red criteria

YES

MOVE TO HIGH ACUITY RESUSCITATION AREA IMMEDIATELY

2

CHECK FOR YELLOW CRITERIA

AIRWAY & BREATHING

- Any swelling/mass of mouth, throat or neck
- Wheezing (no red criteria)

CIRCULATION

- Unable to feed or drink
- Vomits everything
- Ongoing diarrhoea
- Dehydration
- Severe pallor (no red criteria)

DISABILITY

- Restless, continuously irritable or lethargy
- Severe pain

OTHER

NO

- Any infant 8 days to 6 months old
- Malnutrition with visible severe wasting OR oedema of both feet
- Trauma/burn (no red criteria)
- Sexual assault
- Known diagnosis requiring urgent surgical intervention
- New rash worsening over hours or peeling (no red criteria)
- Exposure requiring time-sensitive prophylaxis (e.g. animal bite)
- Pregnancy (no red criteria)
- Headache (no red criteria)



Patients with high-risk vital signs or clinical concern need up-triage or immediate review by supervising clinician



NO

3 CHE

CHECK FOR HIGH-RISK VITAL SIGNS

Temp $<36^{\circ}$ or $>39^{\circ}$

SpO2 < 92%

AVPU other than A

RR	< 1 year	1-4 years	5-12 years
High	50	40	30
Low	25	20	10
HR	< 1 year	1-4 years	5-12 years
High	180	160	140
Low	< 90	< 80	< 70

NO

YES

MOVE TO CLINICAL TREATMENT AREA

MOVE TO LOW ACUITY OR WAITING AREA