



Table 1. Fitted trends in current tobacco use among persons aged 15 years and older

Table 1: Fitted trends in current tobacco use among persons aged 15 years and older										
Year	CURRENT TOBACCO USE (%)									
	Men			Women			Both sexes			
	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Estimated no. of current tobacco users
2000	47.2	61.3	73.8	21.0	31.9	42.8	34.4	46.9	58.6	38,729,600
2005	48.2	60.4	72.3	19.3	26.0	33.7	34.0	43.5	53.3	40,768,100
2010	49.3	59.8	71.2	16.3	21.4	26.6	32.9	40.7	49.0	41,899,500
2015	47.5	59.5	70.0	13.5	17.9	23.4	30.6	38.8	46.8	44,059,900
2020	47.1	59.0	73.4	9.9	15.1	20.6	28.5	37.1	47.1	46,251,600
2025	45.2	58.6	75.8	7.7	12.9	19.1	26.5	35.8	47.5	48,171,000
30% relative reduction target		41.9			15.0			28.5		

Men - current tobacco use (%)



Women - current tobacco use (%)



Both sexes - current tobacco use (%)






Table 2. Fitted trends in current tobacco smoking among persons aged 15 years and older

Year	CURRENT TOBACCO SMOKING (%)										Estimated no. of current tobacco smokers
	Men			Women			Both sexes				
	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Lower 95% credible limit	Point estimate	Upper 95% credible limit		
2000	42.8	53.4	64.7	3.2	5.0	6.9	23.4	29.7	36.4	24,557,100	
2005	41.0	49.3	58.1	2.2	3.1	4.2	22.0	26.6	31.6	24,942,600	
2010	38.2	45.5	53.5	1.4	1.9	2.5	19.9	23.9	28.2	24,547,200	
2015	34.1	42.0	50.6	0.8	1.2	1.7	17.6	21.7	26.3	24,658,000	
2020	30.1	38.9	47.1	0.5	0.8	1.2	15.3	19.9	24.2	24,824,700	
2025	26.4	36.1	45.1	0.3	0.5	0.8	13.4	18.3	23.0	24,644,000	

Men - current tobacco smoking (%)



Women - current tobacco smoking (%)



Both sexes - current tobacco smoking (%)






Table 3. Fitted trends in current cigarette smoking among persons aged 15 years and older

Table 3.11a: Trends in current cigarette smoking among persons aged 25 years and older										
Year	CURRENT CIGARETTE SMOKING (%)									
	Men			Women			Both sexes			Estimated no. of current cigarette smokers
	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Lower 95% credible limit	Point estimate	Upper 95% credible limit	Lower 95% credible limit	Point estimate	Upper 95% credible limit	
2000	35.8	46.0	57.5	1.5	3.0	4.9	19.0	25.0	31.7	20,628,000
2005	32.6	42.4	51.8	1.0	1.9	2.9	17.1	22.5	27.8	21,044,300
2010	30.1	39.0	47.2	0.6	1.2	1.8	15.5	20.2	24.7	20,800,900
2015	28.4	36.0	44.9	0.3	0.7	1.1	14.4	18.5	23.1	20,967,100
2020	24.8	33.2	42.4	0.2	0.5	0.7	12.5	16.9	21.6	21,059,100
2025	22.3	30.9	40.7	0.1	0.3	0.5	11.2	15.6	20.6	21,034,500

Men - current cigarette smoking (%)



Women - current cigarette smoking (%)



Both sexes - current cigarette smoking (%)





Table 4. Fitted age-specific rates of current tobacco use, 2000, 2010, 2018 and 2025

Age (years)	Men				Women				Both sexes			
	2000	2010	2018	2025	2000	2010	2018	2025	2000	2010	2018	2025
15–24	41.5	39.8	38.7	37.2	17.9	11.8	8.3	6.3	29.9	26.0	23.8	22.1
25–39	65.4	62.3	60.8	58.9	24.7	16.2	11.6	8.8	45.6	39.1	36.0	33.8
40–54	80.0	76.7	74.3	72.0	42.4	27.6	19.9	15.0	62.4	53.1	47.4	43.2
55–69	75.1	71.2	69.4	67.4	64.2	42.6	29.9	22.5	69.9	57.6	50.7	45.6
≥ 70	61.3	58.2	55.8	53.8	71.4	46.5	33.2	24.8	66.2	52.5	44.4	39.0

Men - age-specific rates of current tobacco use (%)



Women - age-specific rates of current tobacco use (%)




Table 5. Nationally representative population-based surveys used to calculate above estimates of tobacco use (most recent six surveys)

Survey name	Survey year	Age (years)	Tobacco use		Tobacco smoking		Cigarette smoking		Smokeless use	
			Men	Women	Men	Women	Men	Women	Men	Women
STEPS Survey	2018	18-69	59.6	28.3	46.6	1.0	26.9	28.1
Global Adult Tobacco Survey	2017	15+	46.0	25.2	36.2	0.8	28.7	0.2	16.2	24.8
Non-Communicable Disease Risk Factor Survey	2013	25+	58.5	32.2	41.4	2.6	34.0	31.2
Non-Communicable Disease Risk Factor Survey	2009-10	25+	70.0	34.4	54.8	1.3	29.4	33.6
Global Adult Tobacco Survey	2009	15+	58.0	28.7	44.7	1.5	26.4	27.9
Demographic and Health Survey	2007	15-54	70.3	21.4	...

Table 6. Nationally representative school-based surveys (most recent six surveys)

Survey name	Survey year	Age (years)	Tobacco use		Tobacco smoking		Cigarette smoking		Smokeless use	
			Boys	Girls	Boys	Girls	Boys	Girls	Boys	Girls
Global School-Based Student Health Survey	2014	13-15	13.2	2.1	10.1	1.5
Global Youth Tobacco Survey	2013	13-15	9.2	2.8	4.0	1.1	3.4	...	5.9	2.0
Global Youth Tobacco Survey	2007	13-15	9.1	5.1	2.9	1.1	5.8	4.2
Global Youth Tobacco Survey (Dhaka)	2004	13-15	5.9	4.7	2.3
...
...

Table 7. National surveillance planning

An effective monitoring system needs at least one nationally representative population-based survey and one school-based survey among adolescents to be completed every five years. Based on surveys already completed, future surveys should be run in or before the years indicated below.

	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030
Population-based survey					X					X		
School-based adolescent survey	X					X					X	

Key messages

WHO estimates that, in 2010, about 41% of Bangladesh's population aged 15 years and older were current tobacco users (approximately 41,899,500 people). If tobacco control efforts continue at the same intensity, WHO projects that in 2025 around 36% of the population aged 15 years and older (approximately 48,171,000 people) will be tobacco users.

WHO estimates that, in 2010, about 60% of men and about 21% of women aged 15 years and older in Bangladesh were using tobacco either daily or occasionally. WHO projects that by 2025 the rate will be approximately 59% for men and 13% for women.

In 2010, the highest rate of smoking among men was seen in the age-group 40–54, and among women in the age-group ≥ 70.

WHO Member States adopted a voluntary global target to reduce use of tobacco (smoking and smokeless combined) by 30% by 2025. If Bangladesh adopts this global target, the results indicate that, based on current trends, Bangladesh is unlikely to achieve the target.