12. CHEMICAL FACT SHEETS

Occurrence	Detected in groundwater and drinking-water, usually at concentrations below 20 µg/l, although levels as high as 440 µg/l have been measured ir well water
TDI	14 µg/kg body weight based on a LOAEL of 71.4 mg/kg body weight per day (100 mg/kg body weight per day adjusted for daily dosing) for changes in haematological parameters in a 13-week study in male rats, with an uncertainty factor of 5000 (100 for interspecies and intraspecies variation, 10 for use of a LOAEL and 5 to reflect limitations of the database, including the limited data on in vivo genotoxicity and use of a subchronic study)
Limit of detection	$0.02\mu g/l$ by purge-and-trap GC with an electrolytic conductivity detector or GC-MS
Treatment performance	1 μg/l should be achievable using GAC
Guideline value derivation	
 allocation to water 	10% of TDI
weight	60 kg adult
consumption	2 litres/day
Assessment date	1998
Principal reference	WHO (2003) 1,2-Dichloropropane (1,2-DCP) in drinking-water

1,2-DCP was evaluated by IARC in 1986 and 1987. The substance was classified in Group 3 (not classifiable as to its carcinogenicity to humans) on the basis of limited evidence for its carcinogenicity in experimental animals and insufficient data with which to evaluate its carcinogenicity in humans. Results from in vitro assays for mutagenicity were mixed. The in vivo studies, which were limited in number and design, were negative. In accordance with the IARC evaluation, the evidence from the long-term carcinogenicity studies in mice and rats was considered limited, and it was concluded that the use of a threshold approach for the toxicological evaluation of 1,2-DCP was appropriate.

1,3-Dichloropropane

1,3-Dichloropropane (CAS No. 142-28-9) has several industrial uses and may be found as a contaminant of soil furnigants containing 1,3-dichloropropene. It is rarely found in water.

Reason for not establishing a guideline value	Available data inadequate to permit derivation of health-based guideline value
Assessment date	1993
Principal reference	WHO (2003) 1,3-Dichloropropane in drinking-water

1,3-Dichloropropane is of low acute toxicity. There is some indication that it may be genotoxic in bacterial systems. No short-term, long-term, reproductive or developmental toxicity data pertinent to exposure via drinking-water could be located in the literature. The available data are considered insufficient to permit recommendation of a guideline value.