

12. CHEMICAL FACT SHEETS

Guideline value	2,4,6-Trichlorophenol: 0.2 mg/l (200 µg/l)
Occurrence	Concentrations of chlorophenols in drinking-water usually less than 1 µg/l
Basis of guideline value derivation	Applying the linearized multistage model to leukaemias in male rats observed in a 2-year feeding study (hepatic tumours found in this study were not used for risk estimation because of the possible role of contaminants in their induction)
Limit of detection	0.5–5 µg/l by formation of pentafluorobenzyl ether derivatives; 0.01 µg/l using GC with ECD
Treatment performance	2,4,6-Trichlorophenol concentrations can be reduced using GAC
Additional comments	The guideline value for 2,4,6-trichlorophenol exceeds its lowest reported taste threshold.
Assessment date	1993
Principal reference	WHO (2003) <i>Chlorophenols in drinking-water</i>
Reason for not establishing guideline values	Available data inadequate to permit derivation of health-based guideline values for 2-chlorophenol and 2,4-dichlorophenol
Assessment date	1993
Principal reference	WHO (2003) <i>Chlorophenols in drinking-water</i>

2-Chlorophenol

Data on the toxicity of 2-chlorophenol are limited. Therefore, no health-based guideline value has been derived.

2,4-Dichlorophenol

Data on the toxicity of 2,4-dichlorophenol are limited. Therefore, no health-based guideline value has been derived.

2,4,6-Trichlorophenol

2,4,6-Trichlorophenol has been reported to induce lymphomas and leukaemias in male rats and hepatic tumours in male and female mice. The compound has not been shown to be mutagenic in the Ames test but has shown weak mutagenic activity in other in vitro and in vivo studies. IARC has classified 2,4,6-trichlorophenol in Group 2B (possibly carcinogenic to humans).

Chloropicrin

Chloropicrin, or trichloronitromethane, is formed by the reaction of chlorine with humic and amino acids and with nitrophenols. Its formation is increased in the presence of nitrates. Limited data from the USA indicate that concentrations in drinking-water are usually less than 5 µg/l.

Reason for not establishing a guideline value	Available data inadequate to permit derivation of health-based guideline value
---	--

Assessment date	1993
Principal reference	WHO (2003) <i>Chloropicrin in drinking-water</i>

Decreased survival and body weights have been reported following long-term oral exposure in laboratory animals. Chloropicrin has been shown to be mutagenic in bacterial tests and in in vitro assays in lymphocytes. Because of the high mortality in a carcinogenesis bioassay and the limited number of end-points examined in the 78-week toxicity study, the available data were considered inadequate to permit the establishment of a guideline value for chloropicrin.

Chlorotoluron

Chlorotoluron (CAS No. 15545-48-9) is a pre emergence or early post emergence herbicide that is slowly biodegradable and mobile in soil. There is only very limited exposure to this compound from food.

Guideline value	0.03 mg/l (30 µg/l)
Occurrence	Detected in drinkingwater at concentrations of less than 1 µg/l
TDI	11.3 µg/kg body weight, derived from a NOAEL of 11.3 mg/kg body weight per day for systemic effects in a 2-year feeding study in mice using an uncertainty factor of 1000 (100 for interspecies and intraspecies variation and 10 for evidence of carcinogenicity)
Limit of detection	0.1 µg/l by separation by reversed-phase HPLC followed by UV and electrochemical detection
Treatment performance	0.1 µg/l should be achievable using GAC
Guideline value derivation	
• allocation to water	10% of TDI
• weight	60 kg adult
• consumption	2 litres/day
Assessment date	1993
Principal reference	WHO (2003) <i>Chlorotoluron in drinking-water</i>

Chlorotoluron is of low toxicity following single, short-term and long-term exposures in experimental animals, but it has been shown to cause an increase in adenomas and carcinomas of the kidneys of male mice given high doses for 2 years. As no carcinogenic effects were reported in a 2-year study in rats, it has been suggested that chlorotoluron has a carcinogenic potential that is both species and sex specific. Chlorotoluron and its metabolites have shown no evidence of genotoxicity.

Chlorpyrifos

Chlorpyrifos (CAS No. 2921-88-2) is a broad-spectrum organophosphorus insecticide used for the control of mosquitoes, flies, various crop pests in soil and on foliage, household pests and aquatic larvae. Although it is not recommended for addition to water for